



CAAPE-5 results for Patient 20190921

Name: Sam Sampler	Interviewer: Test Clinician	Interview completed on: 09/23/2019
Facility: Acme HQ	Date of Birth: 04/01/1997	Age when interviewed: 22
Sex: Male	Ethnic Background: Caucasian / White	Highest degree earned: No high school diploma earned
Marital status: Never married	Current employment status: Unemployed	Primary job type: Laborer / unskilled worker
Typical income: \$10,001 to \$20,000		

Diagnostic Indications

The following are diagnostic indications based on the acknowledgement of the respondent. These indications should be reviewed by a properly credentialed professional who can make the appropriate diagnosis based on CAAPE-5 findings and any other relevant information.

SUBSTANCE USE DISORDER INDICATIONS

A **severe substance use disorder** is indicated for **alcohol** by **6** positive DSM-5 diagnostic criteria. Based on indications of the criteria listed in the substance use disorder item summary table

A **severe substance use disorder** is indicated for **amphetamines** by **8** positive DSM-5 diagnostic criteria. Based on indications of the criteria listed in the substance use disorder item summary table

A **severe substance use disorder** is indicated for **heroin** by **8** positive DSM-5 diagnostic criteria. Based on indications of the criteria listed in the substance use disorder item summary table

MENTAL HEALTH DISORDER INDICATIONS

Major Depressive Episode	There are indications of depressive episodes, but they do not appear to reach diagnostic criteria. Further assessment may be indicated.
Manic Episode	There are no indications of a manic episode.
Bipolar Disorder	There are no indications of a bipolar disorder.
① Panic Disorder	The individual reports repeated panic attacks consistent with a diagnosis of panic disorder.

MENTAL HEALTH DISORDER INDICATIONS

①	Posttraumatic Stress	Findings from the interview strongly indicate a probable diagnosis of PTSD.
	Anxiety and Phobias	The individual did not report any indications of potentially problematic anxiety.
	Obsessions/Compulsions	The individual reports no indication of obsessive-compulsive behaviors.
	Psychosis	The individual manifests no indications of psychosis.
①	Antisocial Personality	The individual reports indications of a conduct disorder as an adolescent beyond the minimum required for a diagnosis.
	Paranoid Personality	No diagnostic criteria indicated.
	Schizoid Personality	No diagnostic criteria indicated.
	Borderline Personality	No diagnostic criteria indicated.
	Dependent Personality	No diagnostic criteria indicated.
	Obsessive-Compulsive Personality	No diagnostic criteria indicated.

ASAM RELATED FINDINGS

Dimension 1 Intoxication/withdrawal considerations

The individual reports use of **Amphetamines in the past 24 hours**. Intoxication and/or potential for withdrawal problems should be evaluated.

The individual reports use of **Alcohol, Heroin within the past week**. Risk for possible withdrawal issues should be considered.

Dimension 5 relapse risk

This individual has the following demographic relapse risk factors:

Under age 25, never married, not a high school graduate, unemployed.

The respondent reported the following clinical risk factors:

The individual reports using multiple substances within the previous week.

The respondent reports repeated injection of substances to get high..

CLINICIAN'S COMMENTS

Patient appeared open and cooperative.

Substance Use Disorder Item Summary Tables

Numbers for each substance indicate the number of times a given event occurred in the last 12 months.

USE IN LARGER AMOUNTS OR OVER LONGER TIME	ALCOHOL	AMPHETAMINES	HEROIN
11 . More time using than intended	1	1	1
17 . Drank/used more than intended	1	1	1
DESIRE/ATTEMPTS TO CUT DOWN/QUIT	ALCOHOL	AMPHETAMINES	HEROIN
13 . Wanted to cut down on drinking/use	0	1	1
18 . Set rules to control drinking/use	0	0	0
19 . Failed to follow rules to control drinking/use	0	0	0
TIME SPENT USING/RECOVERING FROM USE	ALCOHOL	AMPHETAMINES	HEROIN
20 . Time spent using > 20 hr. per week	25	25	25
21 . Planning activities around drinking/use	0	0	0
22 . Intoxicated or high more than a day	0	0	0
CRAVING OR STRONG DESIRE/URGE TO USE	ALCOHOL	AMPHETAMINES	HEROIN
15 . Preoccupation with drinking/use	1	1	1
23 . Craving	1	1	1
24 . Desire to drink/use too strong to resist	0	1	1
FAILURE TO FULFILL ROLE OBLIGATIONS	ALCOHOL	AMPHETAMINES	HEROIN
12 . Neglect of responsibilities	1	1	1
25 . Missed work or school because of use	0	1	1
26 . Work or school problems because of use	0	1	1
27 . Financial problems related to drinking/use	0	0	0
SOCIAL OR INTERPERSONAL PROBLEMS	ALCOHOL	AMPHETAMINES	HEROIN
14 . Objections to drinking/use by others	0	1	1
28 . Violence or hit someone while drunk/high	0	0	0
29 . Drinking/use harmed a relationship	0	1	0
30 . Conflicts related to drinking/use	0	1	0
SACRIFICING ACTIVITIES BECAUSE OF USE	ALCOHOL	AMPHETAMINES	HEROIN
31 . Skipped family/social functions due to use	0	0	0
32 . Given up activities to use	0	0	0
33 . Missed opportunities because of drinking/use	0	0	0
USE THAT IS DANGEROUS/DANGEROUS TO BE IMPAIRED	ALCOHOL	AMPHETAMINES	HEROIN
34 . Injection of a substance to get high	--	1	1
35 . Driven any motor vehicle while impaired	1	0	1
36 . Done risky things while drinking/using/impaired	0	0	0
MEDICAL/PSYCHOLOGICAL CONTRAINDICATION	ALCOHOL	AMPHETAMINES	HEROIN
37 . Med. problem possibly caused by drinking/use	0	0	0
38 . Use despite possible exacerbation of med. problem	0	0	0
39.A . Amnesic periods due to drinking	0	--	--
39.B . Amnesic periods related to use	--	0	0
40 . Drinking/use despite an emotional problem	0	0	0

TOLERANCE	ALCOHOL	AMPHETAMINES	HEROIN
41 . Drink at least 5 drinks per occasion	3	--	--
42 . Drank equivalent of a fifth of spirits	0	--	--
43 . Can drink/use more than before with same effect	1	1	1
44 . Same amount produces less effect	1	1	1
WITHDRAWAL	ALCOHOL	AMPHETAMINES	HEROIN
45 . Withdrawal signs/symptoms	0	0	0
46 . Use to relieve withdrawal	0	0	0
NUMBER OF POSITIVE CRITERIA	6	8	8
2-3 positive criteria = mild			
4-5 positive criteria = moderate			
6 or more positive criteria = severe substance use disorder			
47 . Legal problems are no longer a diagnostic criterion	0	0	0
16 . Use to relieve emotional distress (possible self-med.)	0	0	0

Mental Health Disorder Item Summary Tables

MAJOR DEPRESSIVE EPISODE	
48 . Felt depressed for 2 weeks or more	✓
49 . Lost interest in activities/pleasures	✓
51 . Sleep disturbance	✓
52.A/B . Weight gain or loss without intent	
53 . Difficulty thinking/concentrating	✓
54 . Low energy/easily fatigued	
55 . Feelings of worthlessness or guilt	
56 . Agitated or slowed movements	
57 . Thoughts of death, dying, or suicide	
Total	4
58 . Not related to alcohol or drugs	✓
50 . Occurrence within past 2 months	
MANIC EPISODE	
59 . Unusually happy - euphoria	
60 . Unusually agitated or irritable	
61 . Excessive energy/little need for sleep	
63 . Easily distractible/hard to keep focused	
64 . Thinking speeded up/racing thoughts	
65 . Unusually talkative or pressured speech	
66 . Grandiosity	
67 . Unusual/irresponsible behaviors	
Total	0
68 . Not associated with use of alcohol/drugs	
62 . Occurrence within past 2 months	
PANIC DISORDER	
69 . Distinct period of fear/discomfort	✓
70 . Number of such periods:	10
71 . Choking / Shortness of breath/smothering	✓
72 . Feeling dizzy, light headed, faint	
73 . Sweating / Shaking or trembling	✓
74 . Nausea or stomach distress / Chest pains / Pounding or racing heart	✓
75 . Fear of going crazy/losing control / Fear of dying	
Total	6
POSTTRAUMATIC STRESS	
76 . Experienced/witnessed traumatic event	✓
77 . Learned of event involving family/friend	
78 . Memories come back into mind	✓
79 . Recurrent distressing dreams	✓
80 . Feels as though event reoccurring	

- 81 . Intense distress when reminded of event ✓
- 82 . Actively avoid thoughts or feelings ✓
- 83 . Avoidance of reminders of the event ✓
- 84 . Cannot remember some parts of the event ✓
- 85 . Withdrawn/less interested in activities ✓
- 86 . Hard to be happy or feel positive ✓
- 87 . Trouble sleeping, concentrating/anger ✓
- 88 . More easily startled ✓

Total 11

ANXIETY AND PHOBIAS

- 89 . Worry without reason
- 90 . Frequent anxiety about possible events
- 91 . Concentration/forgetting due to anxiety
- 92 . Nervous, keyed up, on edge
- 93 . Fear of public places or open spaces
- 94 . Avoidance interferes with life

Total 0

OBSESSIONS/COMPULSIONS

- 95 . Repeated ideas, thoughts, impulses
- 96 . Actions to neutralize thoughts/impulses
- 97 . Spend much time to overcome thoughts
- 98 . Thoughts or activities interfere with life
- 99 . Repetitive behaviors to relieve anxiety

Total 0

PSYCHOSIS

- 141 . Auditory hallucinations
- 142 . Olfactory or tactile hallucinations
- 143 . Visual hallucinations
- 146 . Delusion of power
- 147 . Unusual speech
- 148 . Unusual behaviors
- 149 . Flat or inappropriate affect
- 150 . Inappropriate motor behavior

Total 0

ANTISOCIAL PERSONALITY

- 110 . Done dangerous things for the fun of it ✓
- 111 . Could have been arrested after age 15 ✓
- 112 . Was arrested since age 15
- 113 . Frequently lied or conned people ✓
- 114 . Done things impulsively ✓
- 115 . Involved in fights
- 116 . Failure to pay bills/debts ✓

Total	5
PARANOID PERSONALITY	
117 . Holds grudges	✓
118 . Concerns about harm or control	
119 . Suspicious of loyalty of family/friends	
120 . Easily offended according to others	✓
Total	2
SCHIZOID PERSONALITY	
121 . Prefer doing things alone	
122 . Don't need/want close friendships	
123 . Lack of strong emotions	✓
Total	1
BORDERLINE PERSONALITY	
124 . Mood can change quickly	
125 . Intense friendships	
126 . Friendships tend not to last long	
127 . Feelings of emptiness or boredom	
128 . Afraid of abandonment	
129 . Impulsive behavior causes problems	✓
Total	1
DEPENDENT PERSONALITY	
130 . Difficulty making decisions	
131 . Lets others make decisions	
132 . Agree with others to avoid offense	✓
133 . Uncomfortable when alone	
134 . Trouble starting things when alone	
135 . Does anything to get reassurance	
Total	1
OBSESSIVE-COMPULSIVE PERSONALITY	
136 . Perfectionist	✓
137 . Keep things with no use	
138 . Pays close attention to details	
139 . Tends to be in charge	
140 . Perceives self as a hard worker	✓
Total	2



TAAD-5 results for Patient 00000002

Name: John Doe	Interviewer: Norman Hoffmann	Interview completed on: 04/06/2020
Date of Birth: 01/10/1970	Age when interviewed: 50	Sex: Male
Ethnic Background: Multiracial / Biracial / Other	Highest degree earned: No high school diploma earned	

KEY STATEMENTS

The individual acknowledged drinking alcohol 2 to 3 days per week.

The individual acknowledged use of drugs about once a week.

Respondent indicates use of drugs to relieve emotional distress.

The individual acknowledged use of **cocaine (powder or crack), sedatives, hallucinogens** in the past 12 months.

SUBSTANCE USE DISORDER INDICATIONS

A **severe** substance abuse disorder is indicated for **Alcohol** by **9** positive DSM-5 diagnostic criteria.

A **severe** substance abuse disorder is indicated for **Drugs** by **7** positive DSM-5 diagnostic criteria.

SUMMARY OF POSITIVE FINDINGS FOR EACH CRITERION	ALCOHOL	DRUGS
Unintended use	1	0
Desire to cut down/stop/Rule failure	1	1
Time spent using	0	0
Craving/compulsion to use	3	3
Role obligation failure	2	0
Social/interpersonal problems	2	2
Sacrifice activities	1	1
Use in hazardous situations	1	0
Use despite med/psych problems	2	1
Tolerance	2	1

SUMMARY OF POSITIVE FINDINGS FOR EACH CRITERION	ALCOHOL	DRUGS
Withdrawal	0	1

Substance Use Summary Tables

INDICATION	ALCOHOL	DRUGS
1/2.B . 4+ days/wk & 5+ drinks		
2.A . Fifth/day		
2.B . 5+ drinks per occasion	✓	
2.C . Tolerance for alcohol	✓	
3 . Drug use 2+ days/week		
4 . Tolerance for drugs		✓
5 . Relieve emotional distress	✓	
6 . Preoccupation	✓	✓
7 . Drank more (than intended)	✓	
8 . Set rules to limit use		✓
9 . Unable to stop	✓	
10 . Withdrawal		
11.A . Drink for hangover		
11.B . Drug use for withdrawal		✓
12 . Physical problems		
13 . Continued use - med.	✓	
14 . Emotional problems	✓	✓
15 . Neglect responsibilities	✓	
16 . Objections to use	✓	✓
17 . Arguments about use	✓	✓
18 . Use damaged relationship		
19 . Missed work/school		
20 . Other work/school problems	✓	
21 . Injury related to use		
22 . Motor vehicle accident		
23 . Driven under influence	✓	
24 . Craving	✓	✓
25 . Use related arrest	✓	✓
26 . Unintended use		
27 . Extended intoxication		
28 . Compulsion to use	✓	✓
29 . 20+ hours/week of use		
30 . Reduced social activities	✓	✓
31 . Unintended time using		
32 . Sacrifice work activities		